

## Medical Policy

### HIV-Associated Lipodystrophy Syndrome

**Policy Number:** 026

	Commercial and Qualified Health Plans	MassHealth
Authorization required	X	X
Notification within 24 hours of service or next business day		
No notification or authorization		

#### Overview

The purpose of this document is to describe the guidelines Mass General Brigham Health Plan utilizes to determine medical necessity for treatment of HIV-associated lipodystrophy syndrome. The treating provider must request prior authorization and provide documentation as outlined in this policy.

Members must meet the general coverage criteria and the criteria for any specific procedure below:

#### Facial

- Dermal filler injections (Sculptra and Radiesse) for facial lipoatrophy
- Autologous fat transplantation

#### Chest

- Liposuction to reduce lipohypertrophy
- Gynecomastia Surgery to reduce lipohypertrophy

#### Abdomen

- Egrifta™ for lipohypertrophy of the abdomen<sup>1</sup>

#### Neck/Upper Back

- Liposuction to reduce lipohypertrophy

#### General Coverage Criteria

Mass General Brigham Health Plan covers medically necessary treatments and procedures noted above for HIV-associated lipodystrophy syndrome when the following are met:

1. The member has a diagnosis of HIV or AIDS; and
2. The medical condition is well documented by clinical notes (photos may be required), that includes a diagnosis of HIV-associated lipodystrophy syndrome, and specifically states that the treatment is necessary for correcting, repairing, or ameliorating the effects of HIV-associated lipodystrophy syndrome; and
3. The requested procedure can be reasonably expected to treat the specific part of the body affected by HIV-associated lipodystrophy syndrome.

#### Specific Criteria for Selected Procedures

##### Face

---

<sup>1</sup> Mass General Brigham Health Plan's retail pharmacy benefit covers Tesamorelin (Egrifta™) for the treatment of HIV-associated lipodystrophy syndrome, specifically lipohypertrophy of the abdomen through the pharmacy program.

Mass General Brigham Health Plan considers Sculptra and Radiesse (the FDA-approved soft tissue fillers dermal injections for facial lipoatrophy due to HIV-associated lipodystrophy syndrome), as well as autologous fat transplantation, to be medically necessary when the general coverage criteria are met. In addition, the provider performing the procedure must be a contracted in-network provider.

Note: Subsequent injections with the above fillers or autologous fat transplantation may be considered medically necessary however prior authorization and clinical notes and documentation from the treating provider are required.

#### **Exclusions**

1. Semipermanent dermal fillers that are not approved by the FDA for the treatment of facial lipoatrophy due to HIV-associated lipodystrophy syndrome.
2. Semipermanent dermal fillers or autologous fat transplantation that is used for any indication other than facial lipoatrophy due to HIV-associated lipodystrophy syndrome.
3. See [General Exclusions](#)

#### **Chest**

##### **Liposuction**

Mass General Brigham Health Plan covers medically necessary liposuction to reduce lipohypertrophy of the chest caused by HIV-associated lipodystrophy syndrome when the member meets the general coverage criteria.

##### **Gynecomastia Surgery**

Mass General Brigham Health Plan covers medically necessary gynecomastia surgery to reduce lipohypertrophy of the chest caused by HIV-associated lipodystrophy syndrome when the member meets the general coverage criteria and when:

1. Liposuction is not indicated to treat HIV-associated lipohypertrophy of the chest.

#### **Exclusions**

1. Breast surgeries or procedures performed outside the treatment of HIV-associated lipodystrophy syndrome solely to enhance a member's appearance or to counteract appearance that occurs through the natural aging process, in the absence of any signs or symptoms of functional abnormalities and/or associated medical complication is considered cosmetic and is not a covered benefit, unless specifically noted in the coverage criteria.
2. See [General Exclusions](#)

#### **Abdomen**

##### **Egrifta™ Injections**

Under Mass General Brigham Health Plan's retail pharmacy benefit, Tesamorelin (Egrifta™) is covered for the treatment of HIV-associated lipodystrophy syndrome, specifically lipohypertrophy of the abdomen when criteria are met and when authorized through the pharmacy program.

#### **Neck/Upper Back**

##### **Liposuction**

Mass General Brigham Health Plan covers medically necessary liposuction to reduce lipohypertrophy of the neck caused by HIV-associated lipodystrophy syndrome when the member meets the general coverage criteria.

#### **Exclusions**

See [General Exclusions](#)



## General Exclusions

1. When the member does not meet the general coverage criteria;
2. For members with a diagnosis of HIV-Associated Lipodystrophy syndrome, coverage of cosmetic surgery and procedures and non-surgical cosmetic dermatology procedures that are solely to enhance a patient's appearance in the absence of any signs or symptoms of functional abnormalities; and/or associated medical complication is considered cosmetic and is not a covered benefit, unless specifically noted otherwise in this coverage criteria. These include but are not limited to the following:
  - a. Hair removal
  - b. Facial implants
  - c. Skin tightening
  - d. Chemical peels
  - e. Laser skin resurfacing
  - f. Thyroid cartilage shaving surgeries
3. Procedures for facial or body augmentation/reduction not associated with HIV-associated lipodystrophy syndrome

## Definitions

**Autologous fat transplantation** — Autologous fat transplantation involves harvesting of a small intact lump of fatty tissue from the abdomen, cervicodorsal area, or elsewhere, that can be processed into small fat "parcels" that are injected by a syringe with local anesthesia.

**HIV-associated Lipodystrophy Syndrome:** A syndrome that occurs in HIV-infected patients in response to some antiretroviral (ARV) drug therapy, characterized by abnormal fat metabolism and deposition. It is not a single syndrome but rather can be composed of three components that present together, or alone: lipoatrophy, lipohypertrophy, and metabolic disturbance (insulin resistance, hypercholesterolemia, and hypertriglyceridemia).

**Egrifta® (Tesamorelin injection):** Egrifta is a self-administered human growth hormone that was approved by the FDA in 2010 for the treatment of lipodystrophy in HIV infected adults. Egrifta induces and maintains a reduction of excess visceral abdominal fat.

**Facial Lipoatrophy:** Facial lipoatrophy is characterized by loss of the buccal and/or temporal fat pads, leading to facial skeletonization with concave cheeks, prominent nasolabial folds, periorbital hollowing, and visible facial musculature. Also referred to as facial lipodystrophy syndrome (LDS), and facial wasting. The two antiretroviral drugs associated with causing HIV Facial Lipoatrophy are Zidovudine (Brand name: Retrovir; a component of Combivir and Trizivir) and Stavudine (Brand name: Zerit).

**Lipoatrophy:** Loss of fat from specific areas of the body, especially from the face, buttocks, and limbs.

**Lipohypertrophy:** Abnormal accumulation of fat, particularly within the abdomen, breast, dorsocervical region (back of neck and shoulders), front of the neck ("horse collar") and subcutaneous tissue (peripheral lipomatosis).

**Soft Tissue Fillers:** Soft tissue fillers, also known as injectable implants, dermal fillers, or wrinkle fillers are medical device implants approved by the FDA for use in helping to create a smoother and/or fuller appearance in the face, including nasolabial folds, cheeks, and lips and for increasing the volume of the back of the hand.

**Radiesse (Calcium hydroxylapatite):** Radiesse is a filler material for correction of moderate to severe facial wrinkles and folds. It was approved by the FDA in 2006 for facial lipoatrophy in people with HIV lipodystrophy. The effects of this material last approximately 18 months.



**Sculptra (Poly-L-lactic acid):** Sculptra is a long lasting filler material that is given in a series of injections over a period of several months. The effects of Sculptra generally become increasingly apparent over time (over a period of several weeks) and its effects may last up to 2 years. In 2004 the FDA approved Sculptra as injectable filler to correct facial lipoatrophy in people with HIV lipodystrophy.

**Related Policies**

- [Breast Surgeries Medical Policy](#)
- [Reconstructive and Cosmetic Procedures](#)

**Codes**

Authorized CPT/HCPCS Codes	Code Description
11950	Subcutaneous injection of filling material (eg, collagen); 1 cc or less
11951	Subcutaneous injection of filling material (eg, collagen); 1.1 to 5.0 cc
11952	Subcutaneous injection of filling material (eg, collagen); 5.1 to 10.0 cc
15876	Suction assisted lipectomy; head and neck
15877	Suction assisted lipectomy; trunk
15878	Suction assisted lipectomy; upper extremity
15879	Suction assisted lipectomy; lower extremity
Q2026	Injection, Radiesse, 0.1 ml
Q2028	Injection, sculptra, 0.5 mg

**Effective**

January 2023: Annual Review. References updated.  
 January 2022: Annual Review. References updated.  
 January 2021: Annual Review. Revised exclusions. References updated.  
 January 2020: Annual Review. References updated.  
 September 2018: Removed exclusion: liposuction for HIV associated lipodystrophy of the abdomen.  
 December 2017: Annual Update.  
 November 2016: Effective date.

**References:**

1. Barton N, Moore R, Prasad K, Evans G. Excisional lipectomy versus liposuction in HIV-associated lipodystrophy. *Arch Plast Surg.* 2021 Nov;48(6):685-690. doi: 10.5999/aps.2020.02285. Epub 2021 Nov 15. PMID: 34818717; PMCID: PMC8627937.
2. Brown, T. Approach to the human immunodeficiency virus-infected patient with lipodystrophy. *J. Clin. Endocrinol. Metab.* 2008;93:2937-2945
3. Carr A, Miller J, Law M, Cooper DA. A syndrome of lipoatrophy, lactic acidemia and liver dysfunction associated with HIV nucleoside analogue therapy: contribution to protease inhibitor-related lipodystrophy syndrome. *AIDS (London, England).* 2000;14(3):F25-F32
4. Carruthers A, Humphrey S, et al. Injectable soft tissue fillers: permanent agents. UpToDate 2016.
5. Centers for Medicare & Medicaid Services (CMS). Decision memo for dermal injections for the treatment of facial lipodystrophy syndrome (FLS) (CAG-00412N). Medicare Coverage Database. Baltimore, MD: CMS; March 23, 2010.
6. Comite SL, Liu FJ, Balasubramanian S, et al. Treatment of HIV-associated facial lipoatrophy with Radiesse FN (Radiesse). *Dermatol Online J* 2004. 10(2):2.



7. Connolly N, Manders E, Riddler S. Suction-assisted lipectomy for lipodystrophy. *AIDS Res Hum Retroviruses* 2004;20:813
8. Donofrio L. Techniques in facial fat grafting. *Aesthet Surg J* 2008;28(6):681-7.
9. Santos APD, Machado DRL, Schwingel A. Anthropometric cutoff points to identify lipodystrophy characteristics in people living with HIV/AIDS: an observational study. *Nutr Hosp.* 2019 Dec 26;36(6):1315-1323. English. doi: 10.20960/nh.02715. PMID: 31657611.
10. Dos Santos AP, Navarro AM, Schwingel A, et al. Lipodystrophy diagnosis in people living with HIV/AIDS: prediction and validation of sex-specific anthropometric models. *BMC Public Health.* 2018 Jun 27;18(1):806. doi: 10.1186/s12889-018-5707-z. PMID: 29945584; PMCID: PMC6020387.
11. Duracinsky M, Leclercq P, Herrmann S, et al. Safety of poly-L-lactic acid (NewFill) in the treatment of facial lipoatrophy: a large observational study among HIV-positive patients. *BMC Infect Dis.* 2014; 1;14:474
12. Falutz J, Allas S, Blot K, et al. Metabolic effects of a growth hormone-releasing factor in patients with HIV. *N Engl J Med* 2007; 257(23):2359-70
13. Falutz J, Mamputu JC, Potvin D, et al. Effectors of tesamorelin (TH9507), a growth hormone-releasing factor analog, in human immunodeficiency virus-infected patients with excess abdominal fat: a pooled analysis of two multicenter, double-blind placebo-controlled phase 3 trials with safety extension data. *J Clin Endocrinol Metab* 2010;95(9):4291-304.
14. Fiorenza CG, Chou SH, Mantzoros CS. Lipodystrophy: pathophysiology and advances in treatment. *Nat Rev Endocrinol Nature Reviews Endocrinology.* 2011;7(3):137-150. doi:10.1038/nrendo.2010.199.
15. Gervasoni C, Ridolfo A, Rovati L, et al. Maintenance of breast size reduction after mastoplasty and switch to a protease inhibitor-sparing regimen in an HIV-positive woman with highly active antiretroviral therapy-associated massive breast enlargement. *AIDS Patient Care STDS* 2002;16(7):307.
16. Gojanovich GS, Shikuma CM, Milne C. Subcutaneous Adipocyte Adenosine Triphosphate Levels in HIV Infected Patients. *AIDS Res Hum Retroviruses.* 2020 Jan;36(1):75-82. doi: 10.1089/AID.2019.0121. Epub 2019 Sep 17. PMID: 31407586; PMCID: PMC6944137.
17. Glesby MJ. Treatment of HIV-associated lipodystrophy. UpToDate. 2020 April 17. [Updated: May 24, 2022.](#)
18. Gold D, Annino D. HIV-associated cervicodorsal lipodystrophy: etiology and management. *Laryngoscope* 2005;115:791.
19. Grunfeld C, Saag M, Cofrancesco J, et al. Regional adipose tissue measured by MRI over 5 years in HIV-infected and control participants indicates persistence of HIV-associated lipoatrophy. *AIDS* 2010;24:1717.
20. Guaraldi G, Fontdevila J, Christensen LH, et al. Surgical correction of HIV-associated facial lipoatrophy. *AIDS* 2011;25:1
21. Massachusetts Session Laws, Chapter 233 of the Acts of 2016, An Act relative to HIV-Associated lipodystrophy syndrome treatment (effective November 8, 2016). Last accessed December 2, 2020
22. Hultman CS, Mcphail LE, Donaldson JH, Wohl DA. Surgical Management of HIV-Associated Lipodystrophy. *Annals of Plastic Surgery.* 2007;58(3):255-263. doi:10.1097/01.sap.0000248128.33465.83.
23. Jagdeo J, Ho D, et al. A systematic review of filler agents for aesthetic treatment of HIV facial lipoatrophy (FLA). *Journal of the American Academy of Dermatology.* 2015;73(6). doi:10.1016/j.jaad.2015.08.040.



24. Lichtenstein KA. Redefining lipodystrophy syndrome: risks and impact on clinical decision making. *J Acquir Immune Defic Syndr*. 2005;39(4):395-400.
25. Martins de Carvalho F, Casal D. et al. HIV-Associated Facial Lipodystrophy: Experience of a Tertiary Referral Center With Fat and Dermis-Fat Compound Graft Transfer. *Eplasty*. 2016 Dec 1;16:e31. PMID: 28123628; PMCID: PMC5259691.
26. Pavlidis L, Spyropoulou GA, Demiri E. Comparing Efficacy and Costs of Four Facial Fillers in Human Immunodeficiency Virus-Associated Lipodystrophy: A Clinical Trial. *Plast Reconstr Surg*. 2018 Oct;142(4):583e-584e
27. Piliero PJ, Hubbard M, King J, et al. Use of ultrasonography-assisted liposuction of the treatment of human immunodeficiency virus-associated enlargement of the dorsocervical fat pad. *Clin Infect Dis* 2003;37:1374
28. Power R, Tat HL, McGill SM, et al. A qualitative study of the psychosocial implications of lipodystrophy syndrome on HIV positive individuals. *Sex Transm Infect* 2003;79:137
29. Roostaeian, J, Jason B, Jarrahy R, et al. Power-assisted liposuction treatment of cervicodorsal fat pad in human immunodeficiency virus-associated lipodystrophy. *Plastic & Reconstructive Surgery* 2008;121(3):135e-136e.
30. Shuck J, Iorio M, Hung R, et al. Autologous fat grafting and injectable dermal fillers for human immunodeficiency virus-associated facial lipodystrophy: a comparison of safety, efficacy, and long-term treatment outcomes. *Plast Reconstr Surg* 2013;131:499
31. Silvers SL, Eviatar JA, Echavez MI, et al. Prospective, open-label, 18-month trial of calcium hydroxylapatite (Radiesse) for facial soft-tissue augmentation in patients with human immunodeficiency virus-associated lipodystrophy: one -year durability. *Plast Reconstr Surg* 2006;118(3 Supple): 34S
32. Soares LR, Casseb JS, Chaba DC. Self-reported lipodystrophy, nutritional, lipemic profile and its impact on the body image of HIV-1-infected persons, with and without antiretroviral therapy. *AIDS Care*. 2020 Oct;32(10):1317-1322. doi: 10.1080/09540121.2019.1687832. Epub 2019 Nov 11. PMID: 31711307.
33. Strauch B, Baum T, Robbins N. Treatment of Human Immunodeficiency Virus–Associated Lipodystrophy with Dermafat Graft Transfer to the Malar Area. *Plastic and Reconstructive Surgery* 2004;113(1):363–70.
34. Sutinen J. Interventions for managing antiretroviral therapy-associated lipodystrophy. *Curr Opin Infect Dis* 2005;18:25
35. Szczerkowska-Dobosz A, Olszewska B, Lemańska M, et al. Acquired facial lipodystrophy: pathogenesis and therapeutic options. *Postepy Dermatol Alergol* 2015; 32(2): 127–133.
36. Thet D, Siritientong T. Antiretroviral Therapy-Associated Metabolic Complications: Review of the Recent Studies. *HIV AIDS (Auckl)*. 2020 Oct 2;12:507-524. doi: 10.2147/HIV.S275314. PMID: 33061662; PMCID: PMC7537841.
37. Tien PC, Schneider MF, Cole SR, et al. Relation of a Stavudine discontinuation to anthropometric changes among HIV-infected women. *J Acquir Immune Defic Syndrom* 2007;44:43
38. Van Rozelaar L, Kadouch JA, Duyndam DA, et al. Semipermanent filler treatment of HIV-positive patients with facial lipodystrophy: long-term follow-up evaluating MR imaging and quality of life. *Aesthet Surg J* 2014; 1;34(1):118-32



39. Verolet CM, Delhumeau-Cartier C, et al. LIPO Group Metabolism. Lipodystrophy among HIV-infected patients: a cross-sectional study on impact on quality of life and mental health disorders. *AIDS Res Ther.* 2015;12:2
40. Wanke C. Epidemiology, clinical manifestations, and diagnosis of HIV-associated lipodystrophy [Internet] 2015 [cited 2016 Oct 11];

