

Cibinqo® (abrocitinib)
Effective 10/01/2022

Plan	<input type="checkbox"/> MassHealth UPPL <input checked="" type="checkbox"/> Commercial/Exchange	Program Type	<input checked="" type="checkbox"/> Prior Authorization <input checked="" type="checkbox"/> Quantity Limit <input type="checkbox"/> Step Therapy
Benefit	<input checked="" type="checkbox"/> Pharmacy Benefit <input type="checkbox"/> Medical Benefit (NLX)		
Specialty Limitations	This medication has been designated specialty and must be filled at a contracted specialty pharmacy.		
Contact Information	Specialty Medications		
	All Plans	Phone: 866-814-5506	Fax: 866-249-6155
	Non-Specialty Medications		
	MassHealth	Phone: 877-433-7643	Fax: 866-255-7569
	Commercial	Phone: 800-294-5979	Fax: 888-836-0730
	Exchange	Phone: 855-582-2022	Fax: 855-245-2134
	Medical Specialty Medications (NLX)		
	All Plans	Phone: 844-345-2803	Fax: 844-851-0882
Exceptions	N/A		

Overview

Cibinqo is indicated for the treatment of adults with refractory, moderate-to-severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies is inadvisable.

Coverage Guidelines

Authorization may be reviewed for members new to AllWays Health Partners who are currently receiving treatment with Cibinqo, excluding when the product is obtained as samples or via manufacturer's patient assistance programs

OR

Authorization may be granted for members when all the following criteria are met, and documentation is provided:

1. Member is 18 years of age or older
2. Documented diagnosis of moderate to severe atopic dermatitis
3. Provider documents affected body surface is greater than or equal to 10% body surface area OR crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
4. Provider documents ONE of the following:
 - a. Inadequate response to treatment with a medium to super-high potency topical corticosteroid (See Appendix) within the past year
 - b. Topical calcineurin inhibitor in the past year
 - c. Topical corticosteroids and topical calcineurin inhibitors are not advisable for the member
 - d. Inadequate response to treatment with Dupixent or use is not advisable for the member.
5. Member will not use the requested medication concomitantly with any other biologic, targeted synthetic DMARD, or potent immunosuppressants such as azathioprine or cyclosporine.

Continuation of Therapy

Reauthorizations requires physician attestation of continuation of therapy and who achieve or maintain a positive clinical response as evidenced by low disease activity (i.e., clear or almost clear skin), or improvement in signs and symptoms of atopic dermatitis (e.g., redness, itching, oozing/crusting). Member cannot use the requested medication concomitantly with any other biologic, targeted synthetic DMARD, or potent immunosuppressants such as azathioprine or cyclosporine.

Limitations

1. Initial approvals will be granted for 4 months
2. Reauthorizations will be granted for 12 months
3. Member has had a documented negative TB test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for persons who are naïve to biologic DMARDs or targeted synthetic DMARDs associated with an increased risk of TB.
 - a. * If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.
4. The following quantity limits apply:

Cibinqo 50mg, 100mg 200mg	30 tablets per 30 days
---------------------------	------------------------

Appendix

Appendix: Relative potency of select topical corticosteroid products

Potency	Drug	Dosage form	Strength
Super-high potency	Augmented betamethasone dipropionate	Ointment, Lotion, Gel	0.05%
	Clobetasol propionate	Cream, Gel, Ointment, Solution, Cream (emollient), Lotion, Shampoo, Foam, Spray	0.05%
	Fluocinonide	Cream	0.1%
	Flurandrenolide	Tape	4 mcg/cm ²
	Halobetasol propionate	Cream, Lotion, Ointment, Foam	0.05%
High potency	Amcinonide	Ointment	0.1%
	Augmented betamethasone dipropionate	Cream	0.05%
	Betamethasone dipropionate	Ointment	0.05%
	Clobetasol propionate	Cream	0.025%
	Desoximetasone	Cream, Ointment, Spray	0.25%
		Gel	0.05%
	Diflorasone diacetate	Ointment, Cream (emollient)	0.05%
	Fluocinonide	Cream, Ointment, Gel, Solution	0.05%
	Halcinonide	Cream, Ointment	0.1%
Halobetasol propionate	Lotion	0.01%	
High potency	Amcinonide	Cream, Lotion	0.1%
	Betamethasone dipropionate	Cream, hydrophilic emollient	0.05%



Potency	Drug	Dosage form	Strength
Medium potency	Betamethasone valerate	Ointment	0.1%
		Foam	0.12%
	Desoximetasone	Cream, Ointment	0.05%
	Diflorasone diacetate	Cream	0.05%
	Fluocinonide	Cream, aqueous emollient	0.05%
	Fluticasone propionate	Ointment	0.005%
	Mometasone furoate	Ointment	0.1%
	Triamcinolone acetonide	Cream, Ointment	0.5%
	Betamethasone dipropionate	Spray	0.05%
	Clocortolone pivalate	Cream	0.1%
	Fluocinolone acetonide	Ointment	0.025%
	Flurandrenolide	Ointment	0.05%
	Hydrocortisone valerate	Ointment	0.2%
	Mometasone furoate	Cream, Lotion, Solution	0.1%
	Triamcinolone acetonide	Cream	0.1%
Ointment		0.05% and 0.1%	
Aerosol Spray		0.2 mg per 2-second spray	

References

1. Cibinqo [package insert]. New York, NY: Pfizer Inc.; January 2022.

Review History

06/22/2022 – Created and reviewed for June P&T. Effective 10/10/2022

