

**Itraconazole solution
Itraconazole capsules
Effective 11/26/2018**

Plan	<input type="checkbox"/> MassHealth UPPL <input checked="" type="checkbox"/> Commercial/Exchange	Program Type	<input checked="" type="checkbox"/> Prior Authorization <input type="checkbox"/> Quantity Limit <input type="checkbox"/> Step Therapy
Benefit	<input checked="" type="checkbox"/> Pharmacy Benefit <input type="checkbox"/> Medical Benefit		
Specialty Limitations	N/A		
Contact Information	Medical and Specialty Medications		
	All Plans	Phone: 877-519-1908	Fax: 855-540-3693
	Non-Specialty Medications		
	All Plans	Phone: 800-711-4555	Fax: 844-403-1029
Exceptions	N/A		

Overview

Itraconazole oral solution and itraconazole capsules are inhibitor of CYP450-dependent synthesis of ergosterol.

Itraconazole oral solution is indicated for the treatment of oral and/or oesophageal candidiasis in HIV-positive or other immunocompromised patients and prophylaxis of fungal infections in neutropenic patients.

Itraconazole capsules are indicated for the treatment of:

1. Superficial dermatomycoses not responding to topical treatment.
2. Fungal keratitis which has failed to respond to topical treatment or where the disease is either progressing rapidly or is immediately sight threatening.
3. Pityriasis versicolor not responding to any other treatment.
4. Vulvovaginal candidiasis not responding to topical treatment.
5. Oral candidiasis in immunocompromised patients.
6. Onychomycosis caused by dermatophytes.
7. Systemic mycoses, only in the following fungal infections:
 - a. Systemic aspergillosis, histoplasmosis, lymphocutaneous/cutaneous sporotrichosis.
 - b. Treatment and maintenance therapy in AIDS patients with disseminated or chronic pulmonary histoplasmosis infection.
 - c. Treatment of oropharyngeal and/or esophageal candidiasis when first line systemic antifungal therapy is inappropriate or has proven ineffective.
 - d. Treatment of non-invasive candidiasis in non-neutropenic patients when first-line systemic antifungal therapy is inappropriate or has proven ineffective. This may be due to underlying pathology, insensitivity of the pathogen or drug toxicity.

Coverage Guidelines

Itraconazole oral solution 10 mg/mL

Authorization may be granted for members when ALL the following criteria are met:

1. Member has required treatment of oropharyngeal candidiasis or esophageal candidiasis.
2. Prescriber has provided documentation of a treatment failure or contraindication with oral fluconazole 200mg daily or greater.
3. Prescriber has provided documentation of ONE of the following:
 - a. Member is 6 years of age or younger.
 - b. Member has an inability to swallow capsules (i.e. dysphagia).

Itraconazole 100mg capsules

Authorization may be granted for members when product is being used for the one of the following conditions **AND** documentation (clinical notes and diagnostic confirmations) have been submitted:

1. Allergic bronchopulmonary aspergillosis
2. Allergic aspergillus sinusitis
3. Treatment of onychomycosis/tinea unguium (toenail and/or fingernail infection) caused by candida species
4. Treatment of onychomycosis/tinea unguium (toenail and/or fingernail infection) caused by non-candida species in a patient who:
 - a. Has tried and failed terbinafine **AND**
 - b. Is immunocompromised, has diabetes or has pain/mobility issues
5. Treatment of vaginal candidiasis in a patient who has tried and failed a topical vaginal antifungal **AND** fluconazole single-dose 150 mg
6. Treatment of tinea versicolor (pityriasis versicolor) in a patient who has tried and failed a topical antifungal agent*, oral fluconazole **AND** oral ketoconazole
 - a. Note: Trial of a topical antifungal agent may be bypassed if the infection involves a large area of the body that which would be difficult to treat with a topical agent*
7. Treatment of tinea capitis or tinea barbae (tinea sycosis) in a patient who has tried and failed oral griseofulvin
8. Treatment of tinea cruris, tinea faciei, or tinea manuum (tinea manus) in a patient who has tried and failed a topical antifungal agent
9. Treatment of tinea corporis (ring worm) in a patient who has tried and failed a topical antifungal agent* **AND** oral fluconazole
 - a. Note: Trial of a topical antifungal agent may be bypassed if the infection involves a large area of the body that which would be difficult to treat with a topical agent*
10. Treatment of tinea imbricata in a patient who has tried and failed **either** oral griseofulvin **OR** terbinafine
11. Treatment of tinea pedis in a patient who either:
 - a. Has tried and failed a topical antifungal agent **OR**
 - b. Has plantar-type or moccasin-type dry chronic tinea pedis
12. Treatment of oropharyngeal candidiasis (oral thrush) in a patient who has tried and failed oral fluconazole at a daily dose \geq 200 mg
13. Prevention of other systemic or superficial fungal infections in an immunocompromised patient who has tried and failed oral fluconazole (if appropriate for indication)
14. Treatment of other systemic or superficial fungal infections in a patient who has tried and failed oral fluconazole (if appropriate for indication)

Limitations

1. For itraconazole solution authorization will be for 450mL (200mg daily) for 21 days.
2. For itraconazole 100mg capsules, the plan allows a maximum of 170 capsules per 12 months.
 - a. The following quantity and timeframe limits apply:



Indication	Approved Quantity/Duration
Allergic bronchopulmonary aspergillosis	60 capsules per month for 180 days
Allergic aspergillus sinusitis	
Prevention/treatment of other systemic or superficial fungal infections in an immunocompromised person	
Onychomycosis (fingernail)	14 capsules per month for 2 months <u>Dosing:</u> 200 mg twice daily for 1 week; repeat 1-week course after 3-week off-time
Onychomycosis (toenail involvement)	Pulse therapy: 14 capsules per month for 3 months <u>Dosing:</u> 200 mg twice daily for 1 week; repeat 1-week course after 3-week off-time
	Standard therapy: 60 capsules per month for 12 consecutive weeks
Tinea versicolor (Pityriasis versicolor)	14 capsules for 7 days
All other indications	60 capsules per month

References

1. Gupta AK, Mays RR, Versteeg SG, et al. Tinea capitis in children: a systematic review of management. *J Eur Acad Dermatol Venereol* 2018;4.
2. SporanoX (itraconazole) oral solution [prescribing information]. Titusville, NJ: Janssen Pharmaceuticals; April 2018
3. SporanoX (itraconazole) capsules [prescribing information]. Titusville, NJ: Janssen Pharmaceuticals; October 2017.
4. Pappas PG, Kauffman CA, Andes DR, et al. Clinical practice guideline for the management of candidiasis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;62(4):e1-e50.[PubMed [26679628](#)]10.1093/cid/civ933
5. Bradley JS, Nelson JD, Kimberlin DK, et al, eds. *Nelson's Pocket Book of Pediatric Antimicrobial Therapy*. 21st ed. American Academy of Pediatrics; 2015
6. Diflucan (fluconazole) tablets and oral suspension [prescribing information]. New York, NY: Pfizer; March 2018
7. Voravutinon V. Oral treatment of tinea corporis and tinea cruris with terbinafine and griseofulvin: a randomized double blind comparative study. *J Med Assoc Thai* 1993; 76:388.
8. Noble SL, Forbes RC, and Stamm PL. Diagnosis and management of common tinea infections. *American Family Physician*. 1998 Jul;58(1):163-74, 177-8.
9. De Backer M, De Vroey C, Lesaffre E, et al. Twelve weeks of continuous oral therapy for toenail onychomycosis caused by dermatophytes: A double-blind comparative trial of terbinafine 250mg/day vs itraconazole 200 mg/day. *Journal of the American Academy of Dermatology*. 1998 May; 38(5 Pt 3):S57-63.
10. Seebacher C, Brasch J, Abeck D, et al. Onychomycosis: Guidelines of the German Society of Dermatology. *Mycoses*. 2007;50:321-327.
11. Roberts DT, Taylor WD, and Boyle J. Guidelines for the treatment of onychomycosis. *British Journal of Dermatology*. 2003;148:402-410.
12. Drake LA, Dinehart SM, Farmer ER, et al. Guidelines of care for superficial mycotic infections of the skin: onychomycosis. *J Am Acad Dermatol*. 1996;34:116-21.



13. Goldstein AO, Goldstein BG. Dermatophyte (tinea) infections. In: Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2014. Available from: <http://www.uptodate.com/utd/index.do>
14. De Berker D. Fungal Nail Disease. *N Engl J Med*. 2009;360:2108-16.
15. Kauffman CA. Treatment of oropharyngeal and esophageal candidiasis. In: Basow DS. UpToDate [database on the internet]. Waltham (MA): UpToDate; 2014. Available at: <http://www.uptodate.com/utd/index/do>
16. Pappas PG, Kauffman CA, Andes D, et al. Clinical Practice Guidelines for the Management of Candidiasis: 2009 Update by the Infectious Diseases Society of America. *Clinical Infectious Diseases*/ 2009;48:503-35.
17. Akhtar S, Masood S, Tabassum S, et al. Efficacy of itraconazole versus fluconazole in vaginal candidiasis. *J Pak Med Assoc*. 2012 Oct;62(10):1049-52.
18. Goldstein BG, Goldstein AO. Tinea Versicolor. In: Basow DS. UpToDate [database on the internet]. Waltham (MA): UpToDate; 2014. Available at: <http://www.uptodate.com/utd/index/do>
19. Gupta AK, Lane D, Paquet M. Systemic review of systemic treatments for tinea versicolor and evidence-based dosing regimen recommendations. *J Cutan Med Surg*. 2014;18(2):79-90.
20. Schwartz RA. Tinea Barbae Treatment & Management. *Medscape*; 2014. Available at: <http://emedicine.medscape.com/article/1091252-treatment>
21. Schuller J, Remme JJ, Rampen FH, et al. Itraconazole in the treatment of tinea pedis and tinea manuum: comparison of two treatment schedules. *Mycoses*. 1998 Dec;41(11-12):515-20.
22. Wingfield AB, Fernandez-Obregon AC, Wignall FS, et al. Treatment of tinea imbricata

Review History

- 12/19/2005 – Reviewed
- 11/27/2006 – Reviewed
- 11/26/2007 – Reviewed and updated
- 11/24/2008 – Reviewed
- 11/23/2009 – Reviewed and updated (oral thrush criteria)
- 11/22/2010 – Reviewed and updated
- 11/28/2011 – Reviewed and updated
- 11/26/2012 – Reviewed and updated
- 12/01/2012 – Reviewed and updated (RxAuth)
- 01/09/2013 – Reviewed and updated (Onmel® plan exclusion; 12/31/12 file)
- 11/25/2013 – Reviewed
- 11/24/2014 – Reviewed and updated
- 11/23/2015 – Reviewed in P&T Meeting
- 11/26/2018 – Reviewed and updated in P&T Meeting
- 03/18/2020 – Reviewed P&T Mtg
- 08/21/2021 – Removed specialty wording and switched Sporanox to itraconazole solution.

