

Savella® (milnacipran) Effective 11/26/2018 ☐ MassHealth UPPL Plan □ Prior Authorization ⊠Commercial/Exchange **Program Type** ☐ Quantity Limit □ Pharmacy Benefit ☐ Step Therapy Benefit ☐ Medical Benefit Specialty N/A Limitations **Medical and Specialty Medications** All Plans Phone: 877-519-1908 Fax: 855-540-3693 Contact Information **Non-Specialty Medications All Plans** Phone: 800-711-4555 Fax: 844-403-1029 **Exceptions** N/A

Overview

Savella® is a selective serotonin and norepinephrine reuptake inhibitor (SNRI) indicated for the management of fibromyalgia.

Coverage Guidelines

Authorization may be granted for members with a diagnosis of fibromyalgia who has been started and stabilized on Savella for at least 30 days excluding when the product is obtained as samples or via manufacturer's patient assistance programs.

OR

Authorization may be granted for members when ALL the following criteria are met:

- 1. Member has a diagnosis of fibromyalgia
- 2. Member has had a documented side effect, allergy, or treatment failure with an SNRI such as a generic venlafaxine product (Effexor® XR caps/tabs, etc.), duloxetine (Cymbalta®), or a desvenlafaxine product [Pristiq®, Khedezla®, etc.]).
- 3. Member has had a documented side effect, allergy, or treatment failure with <u>at least one</u> agent from two <u>different</u> categories including:
 - a. Tricyclic antidepressants (e.g., amitriptyline, doxepin, desipramine, imipramine, etc.)
 - b. SSRI's (e.g., citalopram, fluoxetine, paroxetine, sertraline, etc.)
 - c. Cyclobenzaprine
 - d. Gabapentin

Limitations

- 1. Initial approvals will be granted for 36 months.
- 2. The following quantity limits apply:

	- -
Savella Titration pack	One time only

Savella tablets	60 tablets per 30 days
-----------------	------------------------

References

- 1. Clauw DJ., Mease P., Palmer RH., Gendreau RM., Wang Y. Milnacipran for the treatment of fibromyalgia in adults: a 15-week, multicenter, randomized, double-blind, placebo-controlled, multiple-dose clinical trial. Clinical Therapeutics. 2008; 30(11):1988-2004.
- 2. Mease PJ, Clauw DJ, Gendreau RM, et al. The efficacy and safety of milnacipran for treatment of fibromyalgia: a randomized, double-blind, placebo-controlled trial. J Rheumatol. 2009; 36:398-409.
- 3. Peterson EL. Fibromyalgia--management of a misunderstood disorder. J Am Acad Nurse Pract. 2007 Jul;19(7):341-8.
- 4. Chakrabarty S, Zoorob R. Fibromyalgia. Am Fam Physician. 2007 Jul 15;76(2):247-54.
- 5. Goldenberg DL, Burckhardt C, Crofford L. Management of fibromyalgia syndrome. JAMA.2004; 292(19):2388-2395.
- 6. Carville SF, Arendt-Nielsen S, Bliddal H, et al. EULAR evidence- based recommendations for the management of fibromyalgia syndrome. Ann Rheum Dis. 2008; 67:536-541.
- 7. Vitton O, Gendreau M, Gendreau J, Kranzler J, Rao SG. A double-blind placebo-controlled trial of milnacipran in the treatment of fibromyalgia. Hum Psychopharmacol Clin Exp. 2004; 19: S27-S35.
- 8. Savella (milnacipran) [prescribing information]. Irvine, CA: Allergan USA Inc; December 2017
- 9. Buckhardt CS et al. Guideline for the management of fibromyalgia syndrome pain in adults and children. Glenview (IL): American Pain Society (APS); 2005. 109p (Clinical practice guideline; no. 4).
- 10. Traynor LM, Thiessen CN, Traynor AP. Pharmacotherapy of fibromyalgia. Am J Health-Syst Pharm. 2011;68(July 15, 2011):1307-1319.
- 11. Arnold LM, Palmer RH, Ma Y. A 3-year, open-label, flexible-dosing study of milnacipran for the treatment of fibromyalgia. Clinical Journal of Pain. 2013;29(12):1021-8.
- 12. Sumpton JE, Moulin DE. Fibromyalgia. Handbook of Clinical Neurology. 2014; 119:513-27.

Review History

11/22/2010 - Reviewed

11/28/2011 - Reviewed and updated

11/26/2012 - Reviewed

11/25/2013 – Reviewed and updated

08/04/2014 – Updated (duloxetine generic; 12/30/2013 file)

11/24/2014 – Reviewed and updated in P&T Meeting

11/27/2017 - Reviewed

11/26/2018 – Reviewed in P&T Meeting

01/22/2019 - Reviewed

01/22/2020 - Reviewed at P&T.

