

Xifaxan 550mg (rifaximin) Effective: 04/01/2025

Plan	☐ MassHealth UPPL ☑Commercial/Exchange	Drawaya Tura	☑ Prior Authorization	
Benefit	☑ Pharmacy Benefit☐ Medical Benefit	Program Type	☐ Quantity Limit☐ Step Therapy	
Specialty Limitations	N/A			
Contact Information	Medical and Specialty Medications			
	All Plans	Phone: 877-519-1908	Fax: 855-540-3693	
	Non-Specialty Medications			
	All Plans	Phone: 800-711-4555	Fax: 844-403-1029	
Exceptions	N/A			

Overview

Xifaxan (rifaximin) is a rifamycin antibacterial indicated for:

- 1. Reduction in risk of overt hepatic encephalopathy (HE) recurrence in adults
- 2. Treatment of irritable bowel syndrome with diarrhea (IBS-D) in adults

Coverage Guidelines

Authorization may be granted for members who are new to the plan within the past 90 days currently receiving treatment with requested medication, excluding when the product is obtained as samples or via manufacturer's patient assistance programs.

OR

Authorization will be granted when all the following criteria has been met:

Hepatic Encephalopathy

- 1. Documented diagnosis of hepatic encephalopathy
- 2. Member is 18 years of age or older
- 3. Documentation has had an inadequate response, adverse reaction, or contraindication to lactulose

Irritable Bowel Syndrome (IBS-D)

- 1. Documented diagnosis of Irritable Bowel Syndrome with Diarrhea (IBS-D)
- 2. Member is 18 years of age or older
- 3. Documentation that member has had an inadequate response or has a contraindication to ONE of the following:
 - a. Loperamide
 - b. Diphenoxylate/atropine
- 4. Documentation that member has had an inadequate response or contraindication to a bile acid sequestrant (e.g., cholestyramine, colestipol, colesevelam)

Limitations

- 1. Approvals will be granted for:
 - a. Hepatic Encephalopathy: 12 months
 - b. IBS w/ Diarrhea: 30 days
- 2. The following quantity limits apply:

Diagnosis	Quantity Limitations	
Hepatic Encephalopathy	550mg twice a day; 60 tablets per 30 days	
	550mg three times a day x 14 days	
IBS w/Diarrhea	42 tablets per 14 days – May be treated up to 2 times with	
	this regimen	

References

- 1. Antibiotics for induction and maintenance of remission in Crohn's disease. Cochrane Database Syst Rev. 2019 Feb 7;2:CD012730. doi: 10.1002/14651858.CD012730.pub2.
- 2. Lembo A, Pimentel M, Rao SS, et al. Repeat treatment with rifaximin is safe and effective in patients with diarrhea-predominant irritable bowel syndrome. *Gastroenterology*. 2016;151(6):1113-1121.
- 3. Vilstrup H, Amodio P, Bajaj J, et al. Hepatic encephalopathy in chronic liver disease: 2014 Practice Guideline by the American Association for the Study of Liver Diseases and the European Association for the Study of the Liver. Hepatology. 2014;60(2):715-735
- 4. Xifaxan (rifaximin) [prescribing information]. Bridgewater, NJ: Salix Pharmaceuticals Inc; October 2023.

Review History

06/19/2019: - Reviewed

07/21/2021 - Reviewed July P&T; references update; no clinical changes

11/16/2022 - Reviewed for Nov P&T. Separated out MH vs Comm/Exch. No clinical changes.

10/11/2023 - Reviewed and Updated for Oct P&T; added approval duration. Effective 1/1/24

01/08/2025- Reviewed and updated for January P&T. Updated criteria to require documentation of diagnoses and previous trials. Updated HE criteria to remove allowance for approval if the member is currently on therapy, as that is addressed in the criteria for members who are new to the Plan. Effective 04/01/2025.

