

**Taltz (ixekizumab)**  
**Effective 08/16/2026**

|                              |                                                                                                      |                                            |                                                                                  |
|------------------------------|------------------------------------------------------------------------------------------------------|--------------------------------------------|----------------------------------------------------------------------------------|
| <b>Plan</b>                  | <input type="checkbox"/> MassHealth UPPL<br><input checked="" type="checkbox"/> Commercial/Exchange  | <b>Program Type</b>                        | <input checked="" type="checkbox"/> Prior Authorization                          |
| <b>Benefit</b>               | <input checked="" type="checkbox"/> Pharmacy Benefit<br><input type="checkbox"/> Medical Benefit     |                                            | <input type="checkbox"/> Quantity Limit<br><input type="checkbox"/> Step Therapy |
| <b>Specialty Limitations</b> | This medication has been designated specialty and must be filled at a contracted specialty pharmacy. |                                            |                                                                                  |
| <b>Contact Information</b>   | <b>Medical Benefit</b><br><b>Pharmacy Benefit</b>                                                    | Phone: 833-895-2611<br>Phone: 800-711-4555 | Fax: 888-656-6671<br>Fax: 844-403-1029                                           |
| <b>Exceptions</b>            | N/A                                                                                                  |                                            |                                                                                  |

### Overview

Taltz (ixekizumab) is a humanized interleukin-17A antagonist indicated for the treatment of:

- Moderate-to-severe plaque psoriasis in patients who are candidates for systemic therapy or phototherapy
- Active psoriatic arthritis
- Active ankylosing spondylitis
- Active non-radiographic axial spondyloarthritis with objective signs of inflammation

### Coverage Guidelines

If member is new to the plan (as evidenced by coverage effective date of less than or equal to 90 days), submission of medical records documenting that the member is currently receiving treatment with the requested drug, excluding when the product is obtained as samples or via manufacturer's patient assistance programs

**OR**

Authorization may be granted if the member meets all the following diagnosis-specific criteria:

#### Moderate to severe plaque psoriasis

1. Diagnosis of moderate to severe plaque psoriasis
2. At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
3. ONE of the following:
  - a. Minimum duration of 4-week trial and failure, contraindication, or intolerance to ONE of the following topical therapies
    1. Corticosteroids (e.g., betamethasone, clobetasol)
    2. Vitamin D analogs (e.g., calcitriol, calcipotriene)
    3. Tazarotene
    4. Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
    5. Anthralin
    6. Coal tar
  - b. Member has severe psoriasis that warrants a systemic biologic DMARD or targeted synthetic therapy as first-line therapy.

### Active psoriatic arthritis (PsA)

1. Diagnosis of active psoriatic arthritis
2. ONE of the following:
  - a. Actively inflamed joints
  - b. Dactylitis
  - c. Enthesitis
  - d. Axial disease
  - e. Active skin and/or nail involvement

### Active ankylosing spondylitis

1. Diagnosis of active ankylosing spondylitis
2. Minimum duration of 1-month trial and failure, contraindication or intolerance to two different non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally tolerated doses.

### Non-Radiographic axial spondyloarthritis (nr-axSpa)

1. Diagnosis of active non-radiographical axial spondyloarthritis
2. Member has objective signs of inflammation (e.g., C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints)
3. Minimum duration of 1-month trial and failure, contraindication or intolerance to two different non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally tolerated doses.

### Continuation of Therapy

Requests for reauthorization will be approved when the following criteria are met:

1. Submission of medical records (e.g., chart notes) documenting an improvement in the member's condition, as evidenced by low disease activity or improvement in signs and symptoms of the condition

### Limitations

1. Initial approvals and reauthorizations will be granted for 24 months
2. The following quantity limits apply:

| Drug Name and Dosage Form | Quantity Limit                  |
|---------------------------|---------------------------------|
| Taltz autoinjector        | 1 autoinjector per 28 days      |
| Taltz prefilled syringe   | 1 prefilled syringe per 28 days |

### References

1. Deodhar A, Strand V, Kay J, Braun J. The term 'non-radiographic axial spondyloarthritis' is much more important to classify than to diagnose patients with axial spondyloarthritis. *Ann Rheum Dis* 2016; 75:791.
2. Griffiths CE, Reich K, Lebwohl M, et al. Comparison of ixekizumab with etanercept or placebo in moderate-to-severe psoriasis (UNCOVER-2 and UNCOVER-3): results from two phases 3 randomised trials. *Lancet*. 2015;386(9993):541-51
3. Kimball AB, Luger T, Gottlieb A, et al. Impact of ixekizumab on psoriasis itch severity and other psoriasis symptoms: Results from 3 phase III psoriasis clinical trials. *J Am Acad Dermatol* 2016; 75:1156
4. Leonardi C, Maari C, Philipp S, et al. Maintenance of skin clearance with ixekizumab treatment of psoriasis: Three-year results from the UNCOVER-3 study. *J Am Acad Dermatol* 2018; 79:824



5. Nash P, Kirkham B, Okada M, et al. Ixekizumab for the treatment of patients with active psoriatic arthritis and an inadequate response to tumour necrosis factor inhibitors: results from the 24-week randomised, double-blind, placebo-controlled period of the SPIRIT-P2 phase 3 trial. *Lancet* 2017; 389:2317.
6. Taltz (ixekizumab) [prescribing information]. Indianapolis, IN: Eli Lilly and Co; August 2024.
7. Weber U, Jurik AG, Lambert RG, Maksymowych WP. Imaging in Spondyloarthritis: Controversies in Recognition of Early Disease. *Curr Rheumatol Rep* 2016; 18:58

### **Review History**

02/20/19 – Reviewed

09/18/19 - Added new indication of AS and updated references

11/20/19 – Added Skyrizi as preferred trial for PS

03/16/2022 – Reviewed and Updated for March P&T; Added Rinvoq and Skyrizi as preferred trial for PsA.

Effective 05/01/2022

9/21/2022 – Reviewed and Updated for Sept P&T. added Rinvoq as preferred agent for ankylosing spondylitis.

Effective 11/1/22.

11/15/2023 – Reviewed and Updated at Nov P&T; For Psoriatic Arthritis: 5% BSA changed to at least 3%.

Removed TB requirement. Updated preferred agents to prior use with 3 of the following agents: Cimzia, Enbrel, Humira or biosimilars, Skyrizi, Stelara, Tremfya, AND Cosentyx. Updated topical therapies. For Psoriatic arthritis:

Updated preferred agents to require previous use of TWO of the following: Cimzia, Enbrel, Humira or biosimilars, Rinvoq, Simponi, Skyrizi, Stelara, Tremfya, Xeljanz/XR, AND Cosentyx, Orencia. Removed requirement of traditional DMARD. Separated out ankylosing spondylitis and non-radiographical axial spondyloarthritis criteria. Removed appendix. Effective 1/1/2024

9/11/2024 – Reviewed and updated at September P&T. Added Rinvoq LQ as a step through treatment option for psoriatic arthritis. Effective 11/1/2024.

10/09/2024 – Reviewed and updated at October P&T. Effective 11/1/2024: updated plaque psoriasis criteria to remove requirement of documentation for conventional therapies. Effective 1/1/2025: Removed biologic step requirements for all indications. Updated reauthorization criteria to require documentation of clinical improvement.

06/11/2025 – Reviewed and updated at June P&T. No Changes. Effective 07/01/2025.

10/08/2025 – Reviewed and updated at October P&T. Minor verbiage updates to trial language for ankylosing spondylitis and nr-axSpA; intent remains the same. Effective 01/01/2026.

03/11/2026 – Reviewed and updated for March P&T. Administrative update - changing verbiage in reauthorization criteria from “documentation is submitted” to “submission of medical records (e.g., chart notes...” and updating language for members who are new to the Plan. Effective 05/01/2026.

04/15/2026 - Updated previous trial language for plaque psoriasis, allowing for bypassing conventional therapy if member has severe psoriasis that warrants a biologic DMARD or targeted synthetic therapy as first-line therapy. Effective 08/16/2026.

