

**Juxtapid® (Iomitapide)**  
Effective 02/01/2022

<b>Plan</b>	<input checked="" type="checkbox"/> MassHealth <input type="checkbox"/> Commercial/Exchange	<b>Program Type</b>	<input checked="" type="checkbox"/> Prior Authorization
<b>Benefit</b>	<input checked="" type="checkbox"/> Pharmacy Benefit <input type="checkbox"/> Medical Benefit (NLX)		<input checked="" type="checkbox"/> Quantity Limit <input type="checkbox"/> Step Therapy
<b>Specialty Limitations</b>	This medication has been designated as a limited distribution specialty product and must be filled at a contracted specialty pharmacy; dispensing is available through Dohmen Life Science Services.		
<b>Contact Information</b>	<b>Specialty Medications</b>		
	All Plans	Phone: 866-814-5506	Fax: 866-249-6155
	<b>Non-Specialty Medications</b>		
	MassHealth	Phone: 877-433-7643	Fax: 866-255-7569
	Commercial	Phone: 800-294-5979	Fax: 888-836-0730
Exchange	Phone: 855-582-2022	Fax: 855-245-2134	
	<b>Medical Specialty Medications (NLX)</b>		
	All Plans	Phone: 844-345-2803	Fax: 844-851-0882
<b>Exceptions</b>	N/A		

### Overview

JUXTAPID is a microsomal triglyceride transfer protein inhibitor indicated as an adjunct to a low-fat diet and other lipid-lowering treatments, including LDL apheresis where available, to reduce low-density lipoprotein cholesterol (LDL-C), total cholesterol (TC), apolipoprotein B (apo B), and non-high-density lipoprotein cholesterol (non-HDL-C) with homozygous familial hypercholesterolemia (HoFH).

### Coverage Guidelines

Authorization may be granted for members new to the plan who are currently receiving treatment with Juxtapid, excluding when the product is obtained as samples or via manufacturer’s patient assistance programs

#### OR

Authorization may be granted for members when ALL the following criteria are met:

1. Member has a diagnosis of homozygous familial hypercholesterolemia (HoFH)
2. Member is ≥ 18 years of age
3. Member is adherent to a low-fat diet (< 20% of energy supplied by dietary fat intake) and will be taking a dietary supplement to prevent nutritional deficiencies
4. Member has had a documented side-effect, allergy, inadequate response, treatment failure, or contraindication to treatment with a high potency HMG Co-A reductase inhibitor (e.g. statin), including atorvastatin or rosuvastatin used in combination with ezetimibe, a fibric acid derivative, and/or cholestyramine
5. Member has had an inadequate response, treatment failure, or has a contraindication to lipid apheresis therapy
6. Member has had an inadequate response, treatment failure, or has a contraindication to Kynamro®\*

\* Note: Needle phobia is not considered an adequate justification for not utilizing Kynamro®



## Limitations

1. Initial authorizations will be approved for 3 months.
2. Reauthorizations will be approved for 12 months.
3. The following quantity limits apply:

Juxtapid	30 capsules per 30 days
----------	-------------------------

## References

1. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al; Writing Committee. 2016 ACC expert consensus decision pathway on the role of non-statin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk: a report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol*. 2016;68(1):92-125.[PubMed 27046161]Famier M, Bruckert E. Severe familial hypercholesterolemia:
2. Yusuf S, Bosch J, Dagenais G, et al. Cholesterol Lowering in Intermediate-Risk Persons without Cardiovascular Disease. *N Engl J Med* 2016; 374:2021
3. Juxtapid (lomitapide) [prescribing information]. Cambridge, MA: Aegerion Pharmaceuticals, Inc; September 2020.
4. Kynamro (mipomersen) [prescribing information]. Cambridge, MA: Genzyme Corporation; March 2019
5. Mortensen MB, Nordestgaard BG. Elevated LDL cholesterol and increased risk of myocardial infarction and atherosclerotic cardiovascular disease in individuals aged 70-100 years: a contemporary primary prevention cohort. *Lancet* 2020; 396:1644.
6. National Heart Lung and Blood Institute. Clinical Practice Guidelines and Reports under Development. Available at:
7. <http://www.nhlbi.nih.gov/guidelines/indevelop.htm> . Accessed December 28, 2012.
8. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology guidelines for management of dyslipidemia and prevention of cardiovascular disease. *Endocr Pract*. 2017;23(suppl 2):1-87.[PubMed 28437620]10.4158/EP171764.APPGL.
9. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2019; 73: e285

## Review History

06/01/2018 – Implemented

02/26/2018 – Reviewed

11/26/2018 – Reviewed

01/22/2020 – Added started and stabilized criteria and removed PCSK9 inhibitor trial

09/22/2021 – Reviewed at September P&T; removed diagnosis and age requirement for new members currently on Juxtapid; references updated. Effective 02/01/2022.

09/21/2022 – Reviewed at Sept P&T; no clinical changes; separated out MH vs. Comm/Exch

## Disclaimer

AllWays Health Partners complies with applicable federal civil rights laws and does not discriminate or exclude people on the basis of race, color, national origin, age, disability, or sex.

