

Complement Inhibitors and Miscellaneous Immunosuppressive Agents

Enjaymo (sutimlimab-jome)
Rystiggo (rozanolixizumab-noli)
Soliris (eculizumab)
Syfovre (pegcetacoplan)
Uplizna (inebilizumab-cdon)
Vyvgart (efgartigimod alfa-fcab)
Vyvgart Hytrulo (efgartigimod alfa and hyaluronidase-qvfc)
Effective 11/17/2025

Plan	☒ MassHealth UPPL ☐ Commercial/Exchange	Program Type	☒ Prior Authorization ☐ Quantity Limit ☐ Step Therapy
Benefit	☐ Pharmacy Benefit ☒ Medical Benefit		
Specialty Limitations	N/A		
Contact Information	Medical and Specialty Medications		
	All Plans	Phone: 877-519-1908	Fax: 855-540-3693
	Non-Specialty Medications		
	All Plans	Phone: 800-711-4555	Fax: 844-403-1029
Notes	Additional agents from this class are available through the pharmacy benefit. Please see the MassHealth Drug List for coverage and criteria.		

Overview

Enjaymo (sutimlimab-jome) is an immunoglobulin G (IgG), subclass 4 monoclonal antibody that is indicated to decrease the need for red blood cell (RBC) transfusion due to hemolysis in adults with cold agglutinin diseases (CAD).

Rystiggo (rozanolixizumab-noli) is a neonatal Fc receptor blocker indicated for the treatment of gMG in adult patients who are anti-AChR or anti-MuSK antibody positive.

Soliris (eculizumab) is indicated for the treatment of atypical hemolytic uremic syndrome (aHUS), generalized myasthenia gravis (MG), neuromyelitis optica spectrum disorder (NMOSD), and paroxysmal nocturnal hemoglobinuria (PNH).

Syfovre (pegcetacoplan) is indicated for the treatment of geographic atrophy (GA) secondary to age-related macular degeneration (AMD).

Uplizna (inebilizumab-cdon) is a CD19-directed cytolytic antibody that is presumed to be involved in CD19 binding. Following binding, inebilizumab-cdon depletes lymphocytes derived from B-cell lineage.

Vyvgart (efgartigimod alfa-fcab) is indicated for the treatment of gMG in adult patients who are anti- AChR antibody positive.

Vyvgart Hytrulo (efgartigimod alfa and hyaluronidase-qvfc) is a subcutaneous product combination of efgartigimod alfa, a human IgG1 antibody fragment marketed for intravenous use as Vyvgart (efgartigimod alfa), and recombinant human hyaluronidase PH20 (rHuPH20), which is Halozyme's ENHANZE® drug delivery technology to increase permeability of the subcutaneous tissue by depolymerizing hyaluronan, facilitating subcutaneous delivery of biologics.

Coverage Guidelines

Authorization may be granted for members new to the plan who are currently receiving treatment with the requested product excluding when the product is obtained as samples or via manufacturer's patient assistance programs.

OR

Authorization may be granted for members when all the following criteria are met:

Enjaymo (sutimlimab-jome)

1. Diagnosis of cold agglutinin disease (CAD)
2. Prescriber is a hematologist or consult notes from specialist are provided
3. Member is ≥18 years of age
4. Hb ≤10 g/dL (dated within the last 60 days)
5. Appropriate dosing

Rystiggo (rozanolixizumab-noli)

Generalized Myasthenia Gravis (MG), AChR-antibody+

1. Diagnosis of gMG
2. Member is ≥18 years of age
3. Member is AChR antibody positive
4. Prescriber is a neurologist or consult notes from a neurology office are provided
5. Inadequate response, adverse reaction, or contraindication to pyridostigmine
6. **ONE** of the following:
 - a. **BOTH** of the following:
 - i. Member has severe disease requiring faster onset medication
 - ii. Inadequate response, adverse reaction or contraindication to IVIG or plasmapheresis with glucocorticoids
 - b. Inadequate response or adverse reaction to **TWO** of the following or contraindication to **ALL** of the following immunosuppressant trials:
 - i. azathioprine
 - ii. cyclosporine
 - iii. glucocorticoids (e.g., prednisone)
 - iv. mycophenolate
 - v. tacrolimus
7. Inadequate response, adverse reaction, or contraindication to Vyvgart (efgartigimod alfa-fcab) or Vyvgart Hytrulo (efgartigimod alfa and hyaluronidase-qvfc)
8. Appropriate dosing

Generalized Myasthenia Gravis, MuSK-antibody+

1. Diagnosis of gMG
2. Member is ≥18 years of age
3. Member is MuSK antibody positive
4. Prescriber is a neurologist or consult notes from a neurology office are provided



5. Inadequate response, adverse reaction, or contraindication to pyridostigmine
6. **ONE** of the following:
 - a. **BOTH** of the following:
 - i. Member has severe disease requiring faster onset medication
 - ii. Inadequate response, adverse reaction or contraindication to IVIG or plasmapheresis with glucocorticoids
 - b. Inadequate response or adverse reaction to **TWO** of the following or contraindication to **ALL** of the following immunosuppressant trials:
 - i. azathioprine
 - ii. cyclosporine
 - iii. glucocorticoids (e.g., prednisone)
 - iv. rituximab
 - v. mycophenolate
 - vi. tacrolimus
7. Appropriate dosing

Soliris (eculizumab)

Atypical hemolytic-uremic syndrome (aHUS)

1. Diagnosis of atypical hemolytic-uremic syndrome (aHUS)
2. Prescriber is a hematologist or nephrologist or consult notes from specialist are provided
3. Appropriate dosing

Generalized Myasthenia Gravis (MG), AChR-antibody+

1. Diagnosis of Generalized Myasthenia Gravis
2. Member is ≥ 6 years of age
3. Member is AChR antibody positive
4. Prescriber is a neurologist or consult notes from a neurology office are provided
5. Inadequate response, adverse reaction, or contraindication to pyridostigmine
6. **ONE** of the following:
 - a. **BOTH** of the following:
 - i. Member has severe disease requiring faster onset medication
 - ii. Inadequate response, adverse reaction or contraindication to IVIG or plasmapheresis with glucocorticoids
 - b. Inadequate response or adverse reaction to **TWO** of the following or contraindication to **ALL** of the following immunosuppressant trials:
 - i. azathioprine
 - ii. cyclosporine
 - iii. glucocorticoids (e.g., prednisone)
 - iv. mycophenolate
 - v. tacrolimus
7. Appropriate dosing

Neuromyelitis optica spectrum disorder (NMOSD)

1. Diagnosis of neuromyelitis optica spectrum disorder
2. Prescriber is a neurologist or consult notes from specialist are provided
3. Documentation of a positive serologic test for anti-aquaporin-4 (AQP4)
4. Member is ≥ 18 years of age
5. Appropriate dosing



Paroxysmal nocturnal hemoglobinuria (PNH)

1. Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH)
2. Prescriber is a hematologist or consult notes from specialist are provided
3. Appropriate dosing
4. Member is ≥ 18 years of age

PLE/CHAPLE (off-label indication)

1. Diagnosis of CD55-deficient protein-losing enteropathy (PLE), or complement hyperactivation, angiopathic thrombosis, and protein-losing enteropathy (CHAPLE) disease
2. Member is ≥ 2 months of age
3. Prescriber is a specialist in rare genetic or hematologic diseases or consult notes from specialist are provided
4. Results from genetic testing confirming a CD55 loss-of-function mutation
5. Appropriate dosing

Syfovre (pegcetacoplan)

1. Diagnosis of geographic atrophy (GA) secondary to age-related macular degeneration (AMD)
2. Prescriber is an ophthalmologist
3. Member is ≥ 50 years of age
4. Absence of choroidal neovascularization (CNV or Wet-AMD) in the treatment eye
5. Normal luminance best corrected visual acuity (BCVA) ≥ 24 letters (20/320 Snellen equivalence)
6. Total GA lesion area ≥ 2.5 and ≤ 17.5 mm², with at least 1 lesion ≥ 1.25 mm² if GA is multifocal.
7. Presence of any pattern of hyperautofluorescence in the junctional zone of GA.
8. Requested dosing is 15mg (0.1 mL) once every 25 days to 60 days

Uplizna (inebilizumab-cdon)

Neuromyelitis optica spectrum disorder (NMOSD)

1. Diagnosis of NMOSD
2. Prescriber is a neurologist or consult notes from specialist are provided
3. Positive serologic test for anti-aquaporin-4 (AQP4)
4. Member is ≥ 18 years of age
5. Appropriate dosing

Immunoglobulin G4-related disease (IgG4-RD)

1. Diagnosis of IgG4-RD
2. Member is ≥ 18 years of age
3. Prescriber is a specialist (e.g., nephrologist, endocrinologist, hepatologist) or consult notes from specialist are provided
4. IgG4-RD affecting $\geq$ two organs/sites at any time in disease history
5. Inadequate response, adverse reaction to ONE or contraindication to ALL glucocorticoids (e.g., prednisone, methylprednisolone)
6. Inadequate response, adverse reaction, or contraindication to rituximab
7. Appropriate dosing

Vyvgart (efgartigimod alfa-fcab)

Vyvgart Hytrulo (efgartigimod alfa and hyaluronidase-qvfc)

1. Diagnosis of Generalized Myasthenia Gravis



2. Member is ≥ 18 years of age
3. Member is AChR antibody positive
4. Prescriber is a neurologist or consult notes from a neurology office are provided
5. Inadequate response, adverse reaction, or contraindication to pyridostigmine
6. **ONE** of the following:
 - a. **BOTH** of the following:
 - i. Member has severe disease requiring faster onset medication
 - ii. Inadequate response, adverse reaction or contraindication to IVIG or plasmapheresis with glucocorticoids
 - b. Inadequate response or adverse reaction to **TWO** of the following or contraindication to **ALL** of the following immunosuppressant trials:
 - i. azathioprine
 - ii. cyclosporine
 - iii. glucocorticoids (e.g., prednisone)
 - iv. mycophenolate
 - v. tacrolimus
7. Appropriate dosing

Vyvgart Hytrulo (efgartigimod alfa and hyaluronidase-qvfc)

1. Diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP)
2. Member is ≥ 18 years of age
3. Prescriber is a neurologist or consult notes from neurologist are provided
4. Appropriate dosing
5. **TWO** of the following:
 - a. Inadequate response, adverse reaction, or contraindication to immune globulin
 - b. Inadequate response, adverse reaction, or contraindication to plasma exchange
 - c. **ONE** of the following:
 - i. Inadequate response or adverse reaction to glucocorticoids (e.g., budesonide, methylprednisolone, prednisone)
 - ii. **BOTH** of the following:
 1. Contraindication to glucocorticoids
 2. Inadequate response or adverse reaction to immunosuppressants (e.g., azathioprine, cyclosporine, cyclophosphamide, mycophenolate mofetil, rituximab)

Continuation of Therapy

aHUS, CIDP, PNH, NMOSD

Reauthorization by physician will infer a positive response to therapy.

AMD (Syfovre):

1. Positive response to therapy
2. Member has not developed nAMD (wet AMD)
3. If requested dosing is $\geq$ every 60 days, prescriber has assessed using less frequent dosing

CAD, Generalized MG (Rystiggo, Soliris):

Reauthorization requires physician documentation of a positive response to therapy.

Generalized MG (Vyvgart):



Reauthorization by physician will infer a positive response to therapy.

PLE/CHAPLE disease

Medical records documenting **ALL** of the following:

1. Improvement or no worsening of clinical symptoms (e.g., abdominal pain, bowel movements, facial and peripheral edema)
2. **ONE** of the following:
 - a. Increase in current serum albumin concentration from baseline serum albumin concentration
 - b. Serum albumin concentration stabilized above lower threshold for normal range (≥ 3.5 g/dL)
3. **ONE** of the following:
 - a. Increase in current serum IgG concentration from baseline serum IgG concentration
 - a. Serum IgG concentration stabilized above lower threshold for age-adjusted normal range

Limitations

1. Initial approvals will be granted for the following:
 - a. aHUS, PNH, NMOSD, PLE/CHAPLE: 1 year
 - b. CIDP (Vyvgart Hytrulo): 3 months
 - c. CAD, Generalized MG, AMD (Syfovre), IgG4-RD (Uplizna): 6 months
2. Reauthorizations will be granted for 1 year.

References

1. Soliris® [package insert]. Cheshire (CT): Alexion Pharmaceuticals, Inc.; 2020 Nov.
2. Ultomiris® [package insert] Boston (MA): Alexion Pharmaceutical Inc; 2022 Apr.
3. Enspryng® [package insert]. South San Francisco (CA): Genentech.; 2020 Aug.
4. Uplizna® [package insert]. Gaithersburg (MD): Viela Bio.; 2021 Jul.
5. Vyvgart® [package insert]. Boston (MA): Argenx US, Inc. 2021 Dec.
6. Enjaymo® [prescribing information]. Waltham (MA): Bioverative USA Inc.; 2022 Feb.
7. Niaudet P, Boyer OG. Overview of hemolytic uremic syndrome in children. In: Mattoo TK (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2021 Dec 7]. Available from: <http://www.utdol.com/utd/index.do>.
8. Noris M, Remuzzi G. Atypical hemolytic-uremic syndrome. N Engl J Med. 2009 Oct; 361(17):1,676-87.
9. Niaudet P, Boyer OG. Complement-mediated hemolytic uremic syndrome. In: Mattoo TK (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2021 Dec 7]. Available from: <http://www.utdol.com/utd/index.do>.
10. Patriquin CJ, Kuo KHM. Eculizumab and Beyond: The Past, Present, and Future of Complement Therapeutics. Transfus Med Rev. 2019 Oct 22. pii: S0887-7963(19)30132-4.
11. Bird SJ. Diagnosis of Myasthenia Gravis. In: Shefner JM (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2022 Apr 26 [cited 2022 Jun 25]. Available from: <http://www.utdol.com/utd/index.do>.
12. PNHSOURCE: Paroxysmal Nocturnal Hemoglobinuria (PNH) [webpage on the Internet]. PNHSOURCE Web site. Cheshire: PNHSOURCE [cited 2020 April 20]. Available from: <http://www.pnhsource.com/hcp/facts-about-pnh>.
13. Brodsky R. Clinical manifestations and diagnosis of paroxysmal nocturnal hemoglobinuria. In: Larson RA (Ed). UpToDate [database on the internet]. Waltham (MA):UpToDate; 2021 [cited 2021 Dec 7]. Available from: <http://www.utdol.com/utd/index.do>.
14. Empaveli® [package insert]. Waltham (MA): Apellis Pharmaceuticals, Inc.; 2021 May.
15. Siegel Rare Neuroimmune Association. Neuromyelitis Optica Spectrum Disorder [webpage on the internet]. Powell (OH): Siegel Rare Neuroimmune Association; 2020 [cited 2020 Nov 1]. Available from:



<https://wearesrna.org/living-with-myelitis/disease-information/neuromyelitis-optica-spectrum-disorder/>

16. Glisson CC. Neuromyelitis optica spectrum disorders. In Gonzalez-Scarano F(Ed). UpToDate [database on the Internet]. Waltham (MA): UpToDate; 2021 [cited 2021 Dec 7]. Available from: <http://www.utdol.com/utd/index.do>.
17. Kitley J, Leite MI, Nakashima I, et al. Prognostic factors and disease course in aquaporin-4 antibody-positive patients with neuromyelitis optica spectrum disorder from the United Kingdom and Japan. *Brain*. 2012 Jun;135(Pt 6):1834-49.
18. Jiao Y, Fryer JP, Lennon VA, Jenkins SM, Quek AM, Smith CY et al. Updated estimate of AQP4-IgG serostatus and disability outcome in neuromyelitis optica. *Neurology*. 2013 Oct;81(14):1197-1204.
19. Soliris® (eculizumab) formulary dossier. Alexion, Data on file.
20. Sherman E, Han MH. Acute and Chronic Management of Neuromyelitis Optica Spectrum Disorder. *Curr Treat Options Neurol*. 2015 Nov;17(11):48. PMID: 26433388.
21. Trebst C, Jarius S, Berthele A. Update on the diagnosis and treatment of neuromyelitis optica: recommendations of the Neuromyelitis Optica Study Group (NEMOS). *J Neurol*. 2014 Jan;261(1):1-16. PMID: 24272588.
22. Kessler RA, Mealy MA, Levy M. Treatment of Neuromyelitis Optica Spectrum Disorder: Acute, Preventive, and Symptomatic. *Curr Treat Options Neurol*. 2016 Jan;18(1):2.
23. Huda S, Whittam D, Bhojak M, Chamberlain J, Noonan C, Jacob A. Neuromyelitis optica spectrum disorders. *Clin Med (Lond)*. 2019 Mar;19(2):169-176.
24. FDA Approvals Roundup: Tavneos, Verzenio, Dextenza [press release on the Internet]. Rockville (MD): Regulatory Affairs Professionals Society (RAPS); 2021 Oct 13 [cited 2021 Dec 3]. Available from: <https://www.raps.org/news-and-articles/news-articles/2021/10/fda-approvals-roundup-tavneos-verzenio-dextenza>
25. FDA approves add-on drug for adults with rare form of blood vessel inflammation [press release on the Internet]. Rockville (MD): Food and Drug Administration (US); 2021 Oct 13 [cited 2021 Dec 3]. Available from: <https://www.fda.gov/drugs/news-events-human-drugs/fda-approves-add-drug-adults-rare-form-blood-vessel-inflammation> Administered for the MassHealth Pharmacy Program 21
26. Geetha D, Jefferson AJ. ANCA-Associated Vasculitis: Core Curriculum 2020. *Am J Kidney Dis*. 2020 Jan;75(1):124-137
27. Falk RJ, Merkel PA, King TE. Granulomatosis with polyangiitis and microscopic polyangiitis: Clinical manifestations and diagnosis. In: Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2021 Nov 29]. Available from: <http://www.utdol.com/utd/index.do>
28. Yates M, Watts R. ANCA-associated vasculitis. *Clin Med (Lond)*. 2017 Feb;17(1):60-64.
29. Merkel PA, Kaplan AA, Falk RJ. Granulomatosis with polyangiitis and microscopic polyangiitis: Induction and maintenance therapy. In: Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2021 [cited 2021 Nov 29]. Available from: <http://www.utdol.com/utd/index.do>
30. Kidney Disease: Improving Global Outcomes (KDIGO) Glomerular Diseases Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Glomerular Diseases. *Kidney Int*. 2021 Oct;100(4S):S1-S276.
31. Chung SA, Langford CA, Maz M, Abril A, Gorelik M, Guyatt G, et al. 2021 American College of Rheumatology/Vasculitis Foundation Guideline for the Management of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. *Arthritis Rheumatol*. 2021 Aug;73(8):1366-1383.
32. Jayne DRW, Merkel PA, Schall TJ, Bekker P. Avacopan for the Treatment of ANCA-Associated Vasculitis. *N Engl J Med*. 2021 Feb 18;384(7):599-609.
33. Brugnara C, Berentsen S. Cold agglutinin disease. In: Mentzer WC, Brodsky RA (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; 2022 [cited 2022 May 10]. Available from <http://www.utdol.com/utd/index.do>.



34. Syfovre™ [package insert]. Waltham (MA): Apellis Pharmaceuticals, Inc.; 2023 Feb.
35. FDA approves Syfovre as the first and only treatment for geographic atrophy, a leading cause of blindness. [press release on the internet]. Waltham (MA): Apellis Pharmaceuticals; 2023 February 17 [cited 2023 May 15]. Available from: <https://investors.apellis.com/news-releases/news-release-details/fda-approves-syfovre-pegcetacoplan-injection-first-and-only>
36. American Academy of Ophthalmology. Age-Related Macular Degeneration Preferred Practice Pattern. Ophthalmology. 2019 Oct;127(1):1-65.
37. Lindblad AS, et al, and AREDS Research Group. Arch Ophthalmol. 2009;127(9):1168-1174
38. Apellis Pharmaceuticals. How Syfovre™ works [webpage on the internet]. Waltham (MA): Apellis pharmaceuticals; March 2023 [cited 2023 May 15]. Available from: <https://syfovre.com/about-syfovre/how-syfovre-works/>.
39. Vyvgart Hytrulo® [package insert on the internet]. Boston (MA): argenx US, Inc.; 2023 Jun.
40. Rystiggo® [package insert on the internet]. Smyrna (GA): UCB, Inc.; 2023.

Review History

02/08/2023 - Reviewed and created for Feb P&T; matched MH UPPL criteria to be in compliance with Masshealth unified formulary requirements. Effective 4/1/23

04/12/23 – Reviewed and updated for P&T. Added GPA/MPA to initial approval durations. Effective 6/5/23.

06/14/23 – Reviewed and updated for P&T. Removed preferred product requirement from Enjaymo, Uplizna, Soliris for requests reviewing under MB. Effective 6/30/23.

09/13/23 – Reviewed and updated for P&T. Enjaymo was updated and now requires members to have received a vaccine against encapsulated bacteria at least two weeks prior to treatment initiation. Due to the Medical Benefit Analysis, a decision was made to update Soliris (eculizumab), Vyvgart (efgartigimod alfa-fcab), Uplizna (inebilizumab-cdon) and Enjaymo (sutimlimab-jome) within this guideline to be managed through medical billing and designated with MB. Effective 10/2/23

2/14/24 – Reviewed and updated for P&T. Added Rystiggo and Vyvgart Hytrulo to policy requiring PA through MB (did not include preferred product requirement). Effective 3/4/24

04/10/24 – Reviewed and updated for P&T. Criteria added for Soliris for off label diagnosis of PLE/CHAPLE. New appendix “Age-Adjusted Serum IgG Concentration Reference Ranges” was created. **Syfovre** (pegcetacoplan 150 mg/mL vial) age criteria updated from ≥ 60 to ≥ 50 years of age. Effective 5/6/24

01/2025 – Reviewed and updated for P&T. Removed Tavneos from MB policy as it is being managed through pharmacy. Removed vaccination requirement throughout policy. Criteria for myasthenia gravis indications now includes severity of disease and trial with IVIG or plasmapheresis with glucocorticoids. Added expanded indication of CIDP for Vyvgart Hytrulo. Effective 2/18/25

10/8/25 – Reviewed and updated for P&T. Updated approval durations of drugs used for generalized myasthenia gravis to 6 months initial. Criteria for gMG for Rystiggo was separated by subtype AChR+ and MuSK+. Age was updated to >6 years of age for Soliris in gMG, AChR-antibody+. New indication of IgG4-RD was added for Uplizna. Effective 11/17/25

